

NEET CHEMISTRY 2018-19 - Chennai

Periodic Test :15

Number of questions: 150

Name: _____

ID No: _____

Test ID : 027

Test date: 04.04.2019

Time: 3HRS

Negative Marks : 4 marks for correct attempt & 1 mark deducted for every wrong attempt.

1. In the electrochemical cell : $ZnSO_4(0.01 M) || CuSO_4(1.0 M) | Cu$, the emf of this Daniell cell is E_1 . When the concentration of $ZnSO_4$ is changed to 1.0 M and that of $CuSO_4$ changed to 0.01 M, the emf changes to E_2 . From the followings, which one is the relationship between E_1 and E_2 ? (Given, $RT/F = 0.059$)

- (a) $E_1 < E_2$
- (b) $E_1 > E_2$
- (c) $E_2 = 0.1 E_1$
- (d) $E_1 = E_2$

2. The molecular conductivity of a 0.5 mol/dm^3 solution of $AgNO_3$ with electrolytic conductivity of $5.76 \times 10^{-3} \text{ S cm}^{-1}$ at 298 K is

- a) $2.88 \text{ S cm}^2/\text{mol}$
- (b) $11.52 \text{ S cm}^2/\text{mol}$
- (c) $0.086 \text{ S cm}^2/\text{mol}$
- (d) $28.8 \text{ S cm}^2/\text{mol}$

3. During the electrolysis of molten sodium chloride, the time required to produce 0.10 mol of chlorine gas using a current of 3 ampere is

- a) 55 mins
- b) 110 mins
- (c) 220 mins
- (d) 330 mins

4. If the E_{cell}^0 for a given reaction has a negative value, which of the following gives the correct relationships for the values of ΔG° and K_{eq} ?

- a) $\Delta G^\circ > 0, K_{\text{eq}} < 1$
- b) $\Delta G^\circ > 0, K_{\text{eq}} > 1$
- (c) $\Delta G^\circ < 0, K_{\text{eq}} > 1$
- (d) $\Delta G^\circ < 0, K_{\text{eq}} < 1$

5. The number of electrons delivered at the cathode during electrolysis by a current of 1 ampere in 60 seconds is (charge on electron = $1.60 \times 10^{-19} \text{ C}$)

- a) 6×10^{23}
- b) 6×10^{20}
- (c) 3.75×10^{20}
- (d) 7.48×10^{23}

6. Zinc can be coated on iron to produce galvanized iron but the reverse is not possible. It is because

- a) zinc is lighter than iron
- b) Zinc has lower melting point than iron
- (c) Zinc has lower negative electrode potential than iron
- (d) zinc has higher negative electrode potential than iron

7. The pressure of H_2 required to make the potential of H_2 electrode zero in pure water at 298 K is

- a) 10^{-10} atm
- b) 10^{-4} atm
- c) 10^{-14} atm
- d) 10^{-12} atm

8. A device that converts energy of combustion of fuels like hydrogen and methane, directly into electrical energy is known as

- a) dynamo
- b) Ni-Cd cell
- c) fuel cell
- d) electrolytic cell

9. When 0.1 mol MnO_4^{2-} is oxidized the quantity of electricity required to completely oxidize MnO_4^{2-} to MnO_4^- is

- a) 96500 C
- b) 2×96500 C
- c) 9650 C
- d) 96.50 C


10. The weight of silver (at. wt. = 108) displaced by a quantity of electricity which displaces 5600 mL of O_2 at STP will be

- (a) 5.4 g
- (b) 10.8 g
- (c) 54.0 g
- (d) 108.0 g



11. At 25°C molar conductance of 0.1 molar aqueous solution of ammonium hydroxide is $9.54 \text{ ohm}^{-1} \text{ cm}^2 \text{ mol}^{-1}$ and at infinite dilution its molar conductance is $238 \text{ ohm}^{-1} \text{ cm}^2 \text{ mol}^{-1}$. The degree of ionisation of ammonium hydroxide at the same concentration and temperature is

- (a) 4.008%
- (b) 40.800%
- (c) 2.080%
- (d) 20.800%

12. A button cell used in watches function as following.



If half cell potentials are



$E^{\circ} = 0.34 \text{ V}$ The cell potential will be

- (a) 0.84 V
- (b) 1.34 V
- (c) 1.10V
- (d) 0.42V

13. A hydrogen gas electrode is made by dipping platinum wire in a solution of HCl of pH = 10 and by passing hydrogen gas around the platinum wire at one atm pressure. The oxidation potential of electrode would be

- (a) 0.118 V
- (b) 1.18 V
- (c) 0.059 V
- (d) 0.59 V

14. Consider the half-cell reduction reaction
 $Mn^{2+} + 2e^- \rightarrow Mn$, $E^\circ = -1.18\text{ V}$
 $Mn^{2+} \rightarrow Mn^{3+} + e^-$, $E^\circ = -1.51\text{ V}$ The E° for the
 reaction $3Mn^{2+} \rightarrow Mn + 2Mn^{3+}$, and possibility of
 the forward reaction are respectively

- (a) - 4.18 V and yes
- (b) + 0.33 V and yes
- (c) + 2.69 V and no
- (d) - 2.69 V and no

15. How many gram of cobalt metal will be
 deposited when a solution of cobalt(II) chloride
 is electrolyzed with a current of 10 amperes for
 109 minutes (1 Faraday = 96,500 C; Atomic
 mass of Co = 59 u)

- (a) 4.0
- (b) 20.0
- (c) 40.0
- (d) 0.66

16.

$\Lambda_m^0(\text{NH}_4\text{OH})$ is equal to _____.

- (i) $\Lambda_m^0(\text{NH}_4\text{OH}) + \Lambda_m^0(\text{NH}_4\text{Cl}) - \Lambda_m^0(\text{HCl})$
- (ii) $\Lambda_m^0(\text{NH}_4\text{Cl}) + \Lambda_m^0(\text{NaOH}) - \Lambda_m^0(\text{NaCl})$
- (iii) $\Lambda_m^0(\text{NH}_4\text{Cl}) + \Lambda_m^0(\text{NaCl}) - \Lambda_m^0(\text{NaOH})$
- (iv) $\Lambda_m^0(\text{NaOH}) + \Lambda_m^0(\text{NaCl}) - \Lambda_m^0(\text{NH}_4\text{Cl})$

17. The cell constant of a conductivity cell

- (a) changes with change of electrolyte.

(b) changes with change of concentration of
 electrolyte. O

(c) changes with temperature of electrolyte.

(d) remains constant for a cell.

18. Molar conductivities (Λ_m) at infinite dilution
 of NaCl, MCl and CH_3COONa are 126.4, 425.9
 and 91.0 $\text{S cm}^2 \text{ mol}^{-1}$ respectively. Λ_m for
 CH_3COOH will be

- (a) 425.5 $\text{S cm}^2 \text{ mol}^{-1}$
- (b) 180.5 $\text{S cm}^2 \text{ mol}^{-1}$
- (c) 290.8 $\text{S cm}^2 \text{ mol}^{-1}$
- (d) 390.5 $\text{S cm}^2 \text{ mol}^{-1}$

19. The Gibb's energy for the decomposition of
 Al_2O_3 at 500°C is as follows $\frac{2}{3}\text{Al}_2\text{O}_3 \rightarrow \frac{4}{3}\text{Al} + \text{O}_2$
 $\Delta_r G = +960\text{ kJ mol}^{-1}$. The potential difference
 needed for the electrolytic reduction of
 aluminium oxide (Al_2O_3) at 500°C is at least

- (a) 4.5V
- (b) 3.0V
- (c) 2.5V
- (d) 5.0 V

20. Standard electrode potential of three metals
 X, Y and Z are -1.2 V, + 0.5 V and -0.3 V
 respectively. The reduction power of these
 metals will be

- (a) $Y > Z > X$
- (b) $Y > X > Z$
- (c) $Z > X > Y$
- (d) $X > Y > Z$

21. The electrode potentials for $\text{Cu}^{2+}_{(\text{aq})} + e^- \rightarrow$
 $\text{Cu}^+_{(\text{aq})}$ and $\text{Cu}^+_{(\text{aq})} + e^- \rightarrow \text{Cu}_{(\text{s})}$ are +0.15 V and
 +0.50 V respectively. The value of $E^\circ_{\text{Cu}^{2+}/\text{Cu}}$ will be

- (a) 0.500 V
- (b) 0.325 V
- (c) 0.650 V
- (d) 0.150 V

22. Standard electrode potential for $\text{Sn}^{4+}/\text{Sn}^{2+}$ couple is + 0.15 V and that for the Cr^{3+}/Cr couple is - 0.74 V. These two couples in their standard state are connected to make a cell. The cell potential will be

- (a) + 1.19 V
- (b) + 0.89 V
- (c) +0.18 V
- (d) + 1.83 V

23. A solution contains Fe^{2+} , Fe^{3+} and I^- ions. This solution was treated with iodine at 35°C . E° for $\text{Fe}^{3+}/\text{Fe}^{2+}$ is + 0.77 V and E° for $\text{I}_2/2\text{I}^- = 0.536$ V. The favourable redox reaction is

- (a) I_2 will be reduced to I^-
- (b) there will be no redox reaction
- (c) I^- will be oxidised to I_2
- (d) Fe^{2+} will be oxidised to Fe^{3+}

24. For the reduction of silver ions with copper metal, the standard cell potential was found to be + 0.46 V at 25°C . The value of standard Gibb's energy, ΔG° will be ($F = 96500 \text{ C mol}^{-1}$)

- (a) - 89.0 kJ
- (b) - 89.0 J
- (c) - 44.5 KJ
- (d) - 98.0 kJ

25. An increase in equivalent conductance of a strong electrolyte with dilution is mainly due to

- (a) increase in ionic mobility of ions
- (b) 100% ionisation of electrolyte at normal dilution
- (c) increase in both i.e., number of ions and ionic mobility of ions.
- (d) increase in number of ions.

26. Which of the following expressions correctly represents the equivalent conductance at infinite dilution of $\text{Al}_2(\text{SO}_4)_3$. Given that $\Lambda_{\text{Al}^{3+}}$ and $\Lambda_{\text{SO}_4^{2-}}$ are the equivalent conductances at infinite dilution of the respective ions?

- (a) $2\Lambda_{\text{Al}^{3+}} + 3\Lambda_{\text{SO}_4^{2-}}$
- (b) $\Lambda_{\text{Al}^{3+}} + \Lambda_{\text{SO}_4^{2-}}$
- (c) $(\Lambda_{\text{SO}_4^{2-}} + \Lambda_{\text{SO}_4^{2-}}) \times 6$
- (d) $1/3 \Lambda_{\text{Al}^{3+}} + 1/2 \Lambda_{\text{SO}_4^{2-}}$

27. Consider the following relations for emf of an electrochemical cell

- (i) EMF of cell = (Oxidation potential of anode) - (Reduction potential of cathode)
- (ii) EMF of cell = (Oxidation potential of anode) + (Reduction potential of cathode)
- (iii) EMF of cell = (Reductional potential of anode) + (Reductional potential of cathode)
- (iv) EMF of cell = (Oxidation potential of anode) - (Oxidation potential of cathode)

Which of the above relations are correct?

- (a) (iii) and (i)
- (b) (i) and (ii)
- (c) (iii) and (iv)
- (d) (ii) and (iv)

28. Given :

- (i) $\text{Cu}^{2+} + 2\text{e}^- \rightarrow \text{Cu}$, $E^\circ = 0.337$ V
- (ii) $\text{Cu}^{2+} + \text{e}^- \rightarrow \text{Cu}^+$, $E^\circ = 0.153$ V

Electrode potential, E° for the reaction, $\text{Cu}^+ + \text{e}^- \rightarrow \text{Cu}$, will be

- (a) 0.90V
- (b) 0.30V
- (c) 0.38 V
- (d) 0.52 V


29. Al_2O_3 is reduced by electrolysis at low potentials and high currents. If 4.0×10^4 amperes of current is passed through molten Al_2O_3 for 6 hours, what mass of aluminium is produced? (Assume 100% current efficiency, at. Mass of Al = 27 g mol^{-1}).

- (a) $8.1 \times 10^4 \text{ g}$
- (b) $2.4 \times 10^5 \text{ g}$
- (c) $1.3 \times 10^4 \text{ g}$
- (d) $9.0 \times 10^3 \text{ g}$

30. The equivalent conductance of M/32 solution of a weak monobasic acid is 8.0 mho cm^2 and at the infinite dilution is 400 mho cm^2 . The dissociation constant of this acid is

- (a) 1.25×10^{-6}
- (b) 6.5×10^{-4}
- (c) 1.25×10^{-4}
- (d) 1.25×10^{-5}

31. On the basis of the following E° values, the strongest oxidizing agent is



- (a) Fe^{3+}
- (b) $[\text{Fe}(\text{CN})_6]^{3-}$
- (c) $[\text{Fe}(\text{CN})_6]^{4-}$
- (d) Fe^{2+}

32. Kohlrausch's law states that at

- (a) Infinite dilution, each ion makes definite contribution to conductance of an electrolyte whatever be the nature of the other ion of the electrolyte
- (b) Infinite dilution, each ion makes definite contribution to equivalent conductance of an electrolyte, whatever be the nature of the other ion of the electrolyte
- (c) Finite dilution, each ion makes definite contribution to equivalent conductance of an electrolyte, whatever be the nature of the other ion of the electrolyte
- (d) Infinite dilution each ion makes definite contribution to equivalent conductance of an electrolyte depending on the nature of the other ion of the electrolyte.

33. Standard free energies of formation (in kJ/mol) at 298 K are -237.2 , -394.4 and 4.2 for $\text{H}_2\text{O}_{(l)}$ and $\text{CO}_{2(g)}$, pentane $_{(g)}$ respectively. The value of E°_{cell} for the pentane-oxygen fuel cell is

- (a) 1.0968 V
- (b) 0.0968 V
- (c) 1.968 V
- (d) 2.0968 V

34. The equilibrium constant of the reaction: $\text{Cu}_{(s)} + 2\text{Ag}^+_{(aq)} \rightarrow \text{Cu}^{2+}_{(aq)} + 2\text{Ag}_{(s)}$; $E^\circ = 0.46 \text{ V}$ at 298 K is

- (a) 2.0×10^{10}
- (b) 4.0×10^{10}
- (c) 4.0×10^{15}
- (d) 2.4×10^{10}

35. The efficiency of a fuel cell is given by

- (a) $\Delta G / \Delta S$
- (b) $\Delta G / \Delta N$
- (c) $\Delta S / \Delta G$
- (d) $\Delta H / \Delta G$

36. A hypothetical electrochemical cell is shown below.

$A | A^+ (xM) || B^+ (yM) | B$ The emf measured is + 0.20 V. The cell reaction is

- (a) $A + B^+ \rightarrow A^+ + B$
- (b) $A^+ + B \rightarrow A + B^+$
- (c) $A^+ + e^- \rightarrow A; B^+ + e^- \rightarrow B$
- (d) the cell reaction cannot be predicted.

37. $E^\circ_{Fe^{2+}/Fe} = -0.441$ V and $E^\circ_{Fe^{3+}/Fe^{2+}} = 0.771$ V, the standard EMF of the reaction $Fe + 2Fe^{3+} \rightarrow 3Fe^{2+}$ will be

- (a) 0.111 V
- (b) 0.330 V
- (c) 1.653 V
- (d) 1.212 V

38. 4.5 g of aluminium (at. mass 27 amu) is deposited at cathode from Al^{3+} solution by a certain quantity of electric charge. The volume of hydrogen produced at STP from H^+ ions in solution by the same quantity of electric charge will be

- (a) 44.8 L
- (b) 22.4 L
- (c) 11.2 L
- (d) 5.6 L

39. The mass of carbon anode consumed (giving only carbon dioxide) in the production of 270 kg of aluminium metal from bauxite by the Hall process is


- (a) 270 kg
- (b) 540 kg
- (c) 90 kg
- (d) 180 kg

40. The standard e.m.f of a galvanic cell involving cell reaction with $n = 2$ is found to be 0.295 V at

25° C. The equilibrium constant of the reaction would be

- (a) 2.0×10^{11}
- (b) 4.0×10^{12}
- (c) 1.0×10^2
- (d) 1.0×10^{10}

41. The e.m.f of Daniell cell at 298 K is E_1 .



When the concentration of $ZnSO_4$ is 1M and that of $CuSO_4$ is 0.01 M, the e.m.f changed to E_2 . What is the relationship between E_1 and E_2 ?

- (a) $E_1 > E_2$
- (b) $E_1 < E_2$
- (c) $E_1 = E_2$
- (d) $E_2 = 0 \neq E_1$

42. On the basis of the information available from the reaction,

$4/3Al + O_2 \rightarrow 2/3Al_2O_3$, $\Delta G = -827$ kJ mol⁻¹ of O_2 , the minimum e.m.f required to carry out an electrolysis of Al_2O_3 is ($F = 96500$ C mol⁻¹)

- (a) 2.14 V
- (b) 4.28 V
- (c) 6.47 V
- (d) 8.56 V

43. In electrolysis of NaCl when Pt electrode is taken then H_2 is liberated at cathode while with Hg cathode it forms sodium amalgam

- (a) Hg is more inert than Pt
- (b) More voltage is required to reduce H^+ at Hg than at Pt
- (c) Na is dissolved in Hg while it does not dissolve in Pt
- (d) Conc. of H^+ ions is larger when Pt electrode is taken.

44. Standard electrode potentials are Fe^{2+}/Fe ; $E^\circ = -0.44 \text{ V}$ and $\text{Fe}^{3+}/\text{Fe}^{2+}$ and Fe blocks are kept together, then

- (a) Fe^{3+} increases
- (b) Fe^{3+} decreases
- (c) $\text{Fe}^{2+}/\text{Fe}^{3+}$ remains unchanged
- (d) Fe^{2+} decreases

45. Equivalent conductances of Ba^{2+} and Cl^- ions are $127 \text{ ohm}^{-1} \text{ cm}^{-1} \text{ eq}^{-1}$ respectively. Equivalent conductance of BaCl_2 at infinite dilution is

- (a) 139.5
- (b) 101.5
- (c) 203
- (d) 279

46. For the disproportionation of copper $2\text{Cu}^+ \rightarrow \text{Cu}^{2+} + \text{Cu}$, E° is (Given E° for $\text{Cu}^{2+}/\text{Cu}^+$ is 0.34 V and E° for Cu^{2+}/Cu is 0.15 V .)

- (a) 0.09 V
- (b) -0.19 V
- (c) 0.38 V
- (d) -0.38

47. The specific conductance of a 0.1 N KCl solution at 23°C is $0.012 \text{ ohm}^{-1} \text{ cm}^{-1}$. The resistance of cell containing the solution at the same (=mature was found to be 55 ohm . The cell constant will be

- (a) 0.918 cm^{-1}
- (b) 0.66 cm^{-1}
- (c) 1.142 cm^{-1}
- (d) 1.12 cm^{-1}

48. For the cell reaction, $\text{Cu}^{2+}(\text{C}_1.\text{aq}) + \text{Zn}(\text{s}) = \text{Zn}^{2+}(\text{C}_2.\text{aq}) + \text{Cu}(\text{s})$ of an electrochemical cell, the change in free energy ΔG at a given temperature is a function of

- (a) $\ln(\text{C}_2)$
- (b) $\ln(\text{C}_2/\text{C}_1)$
- (c) $\ln(\text{C}_1)$
- (d) $\ln(\text{C}_1 + \text{C}_2)$

49. E° for the cell, $\text{Zn} | \text{Zn}^{2+}(\text{aq}) || \text{Cu}^{2+}(\text{aq}) | \text{Cu}$ is 1.10 V at 25°C , the equilibrium constant for the reaction $\text{Zn} + \text{Cu}^{2+}(\text{aq}) \rightleftharpoons \text{Cu} + \text{Zn}^{2+}(\text{aq})$ is the order of

- (a) 10^{+8}
- (b) 10^{+17}
- (c) 10^{-28}
- (d) 10^{-37}

50. The molar conductances of NaCl, HCl and CH_3COONa at infinite dilution are 126.45 , 426.16 and $91 \text{ ohm}^{-1} \text{ cm}^2 \text{ mol}^{-1}$ respectively. The molar conductance of CH_3COOH at infinite dilution (A_{m^∞}) is

- (a) $698.28 \text{ ohm}^{-1} \text{ cm}^2 \text{ mol}^{-1}$
- (b) $540.48 \text{ ohm}^{-1} \text{ cm}^2 \text{ mol}^{-1}$
- (c) $201.28 \text{ ohm}^{-1} \text{ cm}^2 \text{ mol}^{-1}$
- (d) $390.71 \text{ ohm}^{-1} \text{ cm}^2 \text{ mol}^{-1}$